arXiv:cond-mat/9712302v1 26 Dec 1997

QUANTUM TUNNELING IN FERRIMAGNETIC NANOPARTICLES COUPLED TO A SPIN BATH: AN EFFECTIVE HAMILTONIAN

I.S. Tupitsyn
Russian Research Center "Kurchatov Institute", 123182 Moscow, Russia.

An effective Hamiltonian, describing quantum tunneling in *ferrimagnetic* nanoparticles which includes interactions between the electronic spins of nanoparticle and microscopic environmental spins (like nuclear spins or paramagnetic impurities), is obtained. Two limiting cases, describing tunneling in antiferromagnetic and ferromagnetic regimes are considered, and criterion for the transition between the two regimes is found. The validity of analytic results is verified by the exact diagonalization method.

I. INTRODUCTION

The quantum tunneling of magnetization (or Neel vector) has been attracting a lot of attention during the last years from both experimental and theoretical point of view since this quantum effect might be seen on macroscopic scale for the total number of spins of order 10^4 and more. Electronic spins in the magnetic grain are coupled to each other by the exchange interaction, forming either a "giant spin" in ferromagnetic case or a "giant Neel vector" in antiferromagnetic case, and can tunnel coherently between the two degenerate states separated by the magnetic anizotropy barrier. The theory of coherent tunneling of the magnetization vector has been developed in many papers [1] (for a review see [2] and refs therein). As to the antiferromagnetic particle, the tunneling effect in this case may be even stronger than in the ferromagnetic one [4] (for a review of modern state of the theory see [3-9]), and of the experiment - [10].) It turns out that in both cases interactions with a spin bath (like nuclear spins or paramagnetic impurities) play a very important role [11,13]. These interactions may be so strong that they can completely suppress the quantum coherence [12]. For the theoretical analysis of the tunneling dynamics one usually introduces an *effective Hamiltonian*, H_{eff}, operating only in the low-energy subspace of the combined system - "giant spin" (or "giant Neel vector") plus the spin bath. This procedure involves truncation of the original problem to the subspace of the two lowest energy levels corresponding to opposite spatial orientations of the order parameter, i.e., all transitions from the lowest doublet to states with exitation energy $E \gtrsim \Omega_o$, where Ω_o characterizes the small oscillation frequency of the order parameter near the classical minima, are ignored in H_{eff}. Earlier H_{eff} for the "giant spin" was derived by the instanton technique in [15]. In this paper I obtain an analogous Hamiltonian for the case of *ferrimagnetic/antiferromagnetic* nanoparticles. To explore the limits of validity of the analytical expressions I use the exact diagonalization technique.

II. HAMILTONIAN AND MODEL

A. STARTING MODEL

Taking advantage of the known fact that Heisenberg model with antiferromagnetic ordering can be reduced in the long-wave limit to the non-linear O(3) σ-model (both in 1D and 2D cases) [3,5,7,9,16], we introduce an effective two-sublattice model with strong exchange interaction between sublattices, the Lagrangian being analogous to that of O(3) σ-model (see, for example, [3,4,8,18]). In addition to the just mentioned works we include coupling between the sublattice spins $\vec{S}_1$ and $\vec{S}_2$ and environmental spins ($\mid \vec{\sigma}_i \mid = \frac{1}{2}$) in a weak magnetic field $\vec{H}_o$. Since coupling to a single environmental spin is typically rather small and can be treated perturbatively, it is sufficient, without loss of generality, to consider here just one such spin coupled, say, to the sublattice $\vec{S}_1$. In this paper we concentrate on the contact hyperfine interaction, assuming that $\vec{\sigma}$ is a nuclear spin. Then, choosing the easy axis/easy plane magnetic anizotropy, we write the starting Hamiltonian as follows:

$$\hat{H} = J\vec{S}_1 \cdot \vec{S}_2 + U(\vec{S}_1) + U(\vec{S}_2) + \frac{\omega_o}{2S_1}\vec{S}_1 \cdot \vec{\sigma}, \;\; U(\vec{S}) = -K_{\parallel}S_z^2 + K_{\perp}S_y^2 - \gamma_e \vec{H}_o \vec{S}, \tag{1}$$

where ω_o is the hyperfine frequency, $K_{\parallel} > 0$ and $K_{\perp} > 0$ are magnetic anizotropy constants, and we assume that J, $K_{\perp} \gg K_{\parallel}$, $\gamma_e S_{1,2} \mid \vec{H}_o \mid \ll 2K_{\parallel}S_{1,2}^2$, $\omega_o \ll 2K_{\parallel}S_{1,2}^2$. (We do not confine ourselves by $J \gg K_{\perp}$, although in practice this limit realizes very often). Assuming $S_1 + S_2 \gg 1$, we will use the *semiclassical* description.

In the low-energy dynamics the dominant contribution to the transition amplitude between the two semiclassical energy minima comes from trajectories with almost *antiparallel* $\vec{S}_1$ and $\vec{S}_2$. That is $\vec{S}_1$ and $\vec{S}_2$ tunnel simultaneously, with only small fluctuations about the comon axis, and the tunneling is described by the Neel vector $\vec{N} = \vec{S}_1 - \vec{S}_2$ with almost constant modulus. As we assume in the general case that $S_1 \neq S_2$, an excess spin $S = S_1 - S_2$ will tunnel together with the Neel vector (ferrimagnetic case).

Let us introduce the spherical angles θ_1, θ_2, ϕ_1, ϕ_2 to describe the orientation of $\vec{S}_1$ and $\vec{S}_2$. We then write $\theta_2 = \pi - \theta_1 - \epsilon_\theta$, $\phi_2 = \pi + \phi_1 + \epsilon_\phi$ with $|\epsilon_\theta|, |\epsilon_\phi| \ll 1$ (see [4]). Also, we choose as the basis set the states characterized by the opposite orientations of Neel vector (along "easy axis" $\vec{z}$), $|\Downarrow\rangle, |\Uparrow\rangle$. Then we can write the transition amplitude as the path integral:

$$\Gamma_{\alpha\beta}(t) = \int_\beta^\alpha \mathcal{D}\{\theta_1, \theta_2, \phi_1, \phi_2\} \exp\left\{ - \int_0^t d\tau [\mathcal{L}_o(\tau) + \delta\mathcal{L}_H(\tau) + \delta\mathcal{L}_\sigma(\tau)] \right\} \;, \tag{2}$$

where $\mathcal{L}_o(\tau)$ - Lagrangian of the non-interacting problem, $\delta\mathcal{L}_H(\tau)$ describing the weak magnetic field contribution and $\delta\mathcal{L}_\sigma(\tau)$ describing the interactions with the nuclear spins (all the Lagrangians are in the Euclidean form and $\alpha, \beta = |\Downarrow\rangle, |\Uparrow\rangle$).

In our problem there are two time scales. Ω_o^{-1} corresponds to the transition time between the two semiclassical minima. The second time scale is defined by the value of the tunneling amplitude Δ_o (the tunneling splitting is equal to $4\Delta_o \cos \pi S$). The relation between the transition amplitude (2) and the non-diagonal part of the effective Hamiltonian is given by [15] (for the real-time axis):

$$\left(H_{eff}^{ND}(\vec{\tau})\right)_{\alpha\beta} = \frac{i}{t}\Gamma^o_{\alpha\beta}(t) \; ; \quad (\alpha \neq \beta \;, \;\; \Omega_o^{-1} \ll t \ll \Delta_o^{-1}) \; . \tag{3}$$

B. SEMICLASSICAL SOLUTION

Let us find first an extremal trajectory for $\theta_1(\tau)$ and $\phi_1(\tau)$ for the non-interacting problem when $|\vec{H}_o| = 0$ and $\omega_o = 0$. The Lagrangian $\mathcal{L}_o$ has the following form:

$$\begin{aligned} \mathcal{L}_o &= JS_1S_2(\sin\theta_1 \sin\theta_2 \cos(\phi_1 - \phi_2) + \cos\theta_1 \cos\theta_2 + 1) + \mathcal{L}^o_{\vec{S}_1} + \mathcal{L}^o_{\vec{S}_2}, \\ \mathcal{L}^o_{\vec{S}} &= -iS\phi\dot{\theta}\sin\theta + K_\parallel S^2 \sin^2\theta + K_\perp S^2 \sin^2\theta \sin^2\phi. \end{aligned} \tag{4}$$

Since $\vec{y}$-axis is a "hard" one, ϕ will weakly fluctuate around $\phi = 0$ or $\phi = \pi$. These values of ϕ correspond to two possible trajectories connecting two semiclassical minima: clockwise and anticlockwise. Performing the Gaussian integration over three "fast" variables (ϕ, ϵ_θ, ϵ_ϕ) we get (omitting all the terms which do not contribute to the equation of motion):

$$\begin{aligned} \mathcal{L}_o(\theta) &= \frac{\mathcal{M}}{2}\dot{\theta}^2 + \frac{S_2^2}{2\tilde{J}}\dot{\phi}^2 \sin^2\theta + \tilde{K}_\parallel \sin^2\theta, \quad \tilde{z} = \tilde{K}_\perp - \frac{2K_\perp^2 S_2^4}{\tilde{J} + 2K_\perp S_2^2}, \\ \mathcal{M} &= \frac{S^2}{2\tilde{z}} + \frac{2SS_2^3K_\perp}{\tilde{z}(\tilde{J} + 2K_\perp S_2^2)} + \frac{S_2^2}{\tilde{J} + 2K_\perp S_2^2} + \frac{2S_2^6K_\perp^2}{\tilde{z}(\tilde{J} + 2K_\perp S_2^2)^2} \;, \end{aligned} \tag{5}$$

where $\tilde{K}_{\perp,\parallel} = K_{\perp,\parallel}(S_1^2 + S_2^2)$ and $\tilde{J} = JS_1S_2$ (below we will ignore indexes at θ and ϕ). This Lagrangian results in the equation of motion $\dot{\theta} = \Omega_o \sin\theta$, the solution of which is:

$$\sin\theta(\tau) = 1/\cosh(\Omega_o\tau), \quad \Omega_o = (2\tilde{K}_\parallel/\mathcal{M})^{1/2} \;, \tag{6}$$

Substituting the extremal trajectory into $\mathcal{L}_o$ and integrating over τ, we get for the Euclidean action:

$$A_o^\eta = A_o + i\eta\pi S, \quad A_o = 4\tilde{K}_\parallel/\Omega_o, \tag{7}$$

where $\eta = \pm$ corresponds to the motion between two semiclassical minima in clockwise or anticlockwise direction and $\eta\pi S$ is the Kramers-Haldane phase.

C. MAGNETIC FIELD

Now we introduce a weak magnetic field ($\gamma_e = 1$):

$$\delta\mathcal{L}_H = \delta\mathcal{L}^H_{\vec{S}_1} + \delta\mathcal{L}^H_{\vec{S}_2}, \;\; \delta\mathcal{L}^H_{\vec{S}} = -S(H_o^x \sin\theta\cos\phi + H_o^y \sin\theta\sin\phi + H_o^z \cos\theta). \tag{8}$$

If we integrate $\mathcal{L}_o + \delta\mathcal{L}_H$ over $\epsilon_\theta, \epsilon_\phi$, with the excess spin S equal to zero and take the limit $J \gg K_\perp$, then we will get the well-known expression for the Lagrangian of Andreev-Marchenko [17] without the gradient term and the term describing weak ferromagnetism of Dzyaloshinskii. But we will proceed in the general form. After integrating (8) over fast variables $\epsilon_\theta, \epsilon_\phi$ and ϕ (keeping only linear in magnetic field terms) we substitute the extremal trajectory (6) and further integrate over τ to get a correction to the action (7):

$$\begin{aligned}\delta A_H^\eta = & -i\frac{\eta\pi S_2^2 H_y}{\tilde{J} + 2K_\perp S_2^2}\left\{1 + \frac{2K_\perp^2 S_2^4}{\tilde{z}(\tilde{J} + 2K_\perp S_2^2)}\right\} - \frac{\eta\pi H_x}{\Omega_o}\left\{\frac{K_\parallel S_2^3}{\tilde{J}} + \frac{K_\perp S_2^5 \Omega_o^2}{\tilde{z}(\tilde{J} + 2K_\perp S_2^2)^2}\right\} \\ & - \frac{\eta\pi S}{\Omega_o}\left\{H_x + i\frac{S\Omega_o}{2\tilde{z}}H_y\right\} - \frac{\eta\pi S\Omega_o S_2^2}{2\tilde{z}(\tilde{J} + 2K_\perp S_2^2)}\left\{H_x + i\frac{4K_\perp S_2}{\Omega_o}H_y\right\} \; .\end{aligned} \tag{9}$$

D. NUCLEAR SPINS

Let us include now coupling between the electronic spins of the nanoparticle and nuclear spins. The correction to the Lagrangian $\mathcal{L}_o$ from hyperfine interaction has a form:

$$\delta\mathcal{L}_\sigma = \frac{\omega_o}{2}\Big(\hat{\sigma}_x \sin\theta_1\cos\phi_1 + \hat{\sigma}_y \sin\theta_1 \sin\phi_1 + \hat{\sigma}_z \cos\theta_1\Big), \tag{10}$$

where $\hat{\sigma}_i$ $(i = x, y, z)$ are Pauli matrixes. Performing the same operations as in the case of magnetic field we get for the correction to the action:

$$\delta A_\sigma^\eta = \frac{\eta\pi\omega_o}{2\Omega_o}\left(\hat{\sigma}_x + i\frac{S_2^3\Omega_o K_\perp}{\tilde{z}(\tilde{J} + 2K_\perp S_2^2)}\hat{\sigma}_y\right) + i\frac{\eta\pi\omega_o S}{4\tilde{z}}\hat{\sigma}_y \; . \tag{11}$$

E. EFFECTIVE HAMILTONIAN

Since $\Omega_o^{-1} \ll t \ll \Delta_o^{-1}$, we can write the non-diagonal part of the effective Hamiltoniah in the form:

$$H_{eff}^{ND} = \frac{i}{t}\left\{\hat{\tau}_- \hat{\Gamma}_{\Downarrow\Uparrow}(t) + H.c.\right\}, \;\; \Gamma_{\Downarrow\Uparrow}(t) = it\Delta_o \sum_{\eta=\pm}\exp\left\{-A^\eta\right\}, \tag{12}$$

where $\hat{\tau}_-$ is the lowering operator in the space of Pauli matrixes and the transition amplitude is (see also [14]):

$$\Delta_o = \frac{\Omega_o}{2}\sqrt{\frac{6}{\pi}A_o}\exp\{-A_o\} \; . \tag{13}$$

We start constructing the effective Hamiltonian from the antiferromagnetic case, when $S_1 = S_2$ and only Neel vector undergoes tunneling. Keeping only linear in H_o and ω_o terms in the action (for the simplicity we set $H_x = 0$) we get:

$$\begin{aligned} A^\eta(0) = A_o(0) - i\eta\psi(0) + \eta\alpha(0)\Big(\hat{\sigma}_x + i\lambda(0)\hat{\sigma}_y\Big), \; \Omega_o(0) = 2S_2\sqrt{K_\parallel(J + K_\perp)}, \\ A_o(0) = 4S_2\lambda(0), \; \psi(0) = \frac{\pi H_y}{J + K_\perp}, \; \alpha(0) = \frac{\pi\omega_o}{2\Omega_o(0)}, \; \lambda(0) = \sqrt{K_\parallel/(J + K_\perp)}. \end{aligned} \tag{14}$$

Substituting (14) into (12), we obtain the non-diagonal part of the effective Hamiltonian:

$$H_{eff}^{ND}(0) = 2\Delta_o(0)\hat{\tau}_- \cos\left[\psi(0) + \alpha(0)\Big(i\hat{\sigma}_x - \lambda(0)\hat{\sigma}_y\Big)\right] + H.c.. \tag{15}$$

We do not present here the diagonal (static) part. The general way of calculating this value one can find in [15].

It is easy to verify that in opposite limiting case, $S_1 \gg S_2$, we will get the effective Hamiltonian describing tunneling of the ferromagnetic-grain magnetization:

$$H^{ND}_{eff}(S_1) = 2\Delta_o(S_1)\hat{\tau}_- \cos\left[\pi S_1 - \psi(S_1) - \alpha(S_1)\Big(i\hat{\sigma}_x - \lambda(S_1)\hat{\sigma}_y\Big)\right] + H.c. \ ,$$

$$\Omega_o(S_1) = 2S_1\sqrt{K_\| K_\perp}, \;\; \alpha(S_1) = \frac{\pi\omega_o}{2\Omega_o(S_1)}, \;\; \psi(S_1) = \frac{\pi H_y}{2K_\perp}, \;\; A_o(S_1) = 2S_1\lambda(S_1), \;\; \lambda(S_1) = \sqrt{\frac{K_\|}{K_\perp}}, \tag{16}$$

where S_1 is understood as the grain's "giant spin".

Finally, in the case of arbitrary value of S (ferrimagnetic) the Neel vector tunnels *together* with the excess spin. In the limit of $\tilde{J} \gg 2K_\perp S_2^2$ the effective Hamiltonian has the following form (as before $H_x = 0$):

$$\begin{aligned} H^{ND}_{eff}(S) &= 2\Delta_o(S)\hat{\tau}_- \cos\left[\pi S - \psi(S) - \alpha(S)\Big(i\hat{\sigma}_x - \lambda(S)\hat{\sigma}_y\Big)\right] + H.c., \\ A_o(S) &= \sqrt{4\frac{K_\|}{K_\perp}\left(S^2 + S_2^2\frac{2\tilde{K}_\perp}{\tilde{J}} + SS_2^3\frac{4K_\perp}{\tilde{J}}\right)}, \;\; \lambda(S) = \frac{\Omega_o(S)}{\tilde{K}_\perp}\left(\frac{S}{2} + \frac{K_\perp S_2^3}{\tilde{J}}\right), \\ \psi(S) &= \frac{\pi A_o(S) H_y}{2\Omega_o(S)}, \;\; \alpha(S) = \frac{\pi\omega_o}{2\Omega_o(S)}, \;\; \Omega_o(S) = \sqrt{\frac{4\tilde{K}_\|\tilde{K}_\perp\tilde{J}}{S^2\tilde{J} + 2\tilde{K}_\perp S_2^2 + 4SS_2^3K_\perp}}. \end{aligned} \tag{17}$$

From the analysis of these three Hamiltonians the criterion (on the value of the excess spin), corresponding to the switching between the ferromagnetic and the antiferromagnetic regimes, follows directly:

$$\begin{aligned} S &\gg S_2\sqrt{\frac{2\tilde{K}_\perp}{\tilde{J} + 2K_\perp S_2^2}} \quad - \;\; ferromagnetic\; regime \\ S &\ll S_2\sqrt{\frac{2\tilde{K}_\perp}{\tilde{J} + 2K_\perp S_2^2}} \quad - \;\; antiferromagnetic\; regime. \end{aligned} \tag{18}$$

III. EXACT DIAGONALIZATION

To verify the obtained expressions we use the exact diagonalization method in application to the starting Hamiltonian (1) with $H_x = 0$. Because of the limited space, we present the numerical results only for the antiferromagnetic case ($S_1 = S_2$). The exact diagonalization procedure is described in detail in [15], and therefore we define here only the values that are subject to the analysis. Expanding (15) into the series up to the linear in ω_o terms we get:

$$H^{ND}_{eff} \approx 2\Delta_o(0)\Big(\cos\psi(0)\cdot\hat{\tau}_x\hat{\sigma}_o - \alpha(0)\sin\psi\cdot\hat{\tau}_y\hat{\sigma}_x + \alpha(0)\lambda(0)\sin\psi\cdot\hat{\tau}_x\hat{\sigma}_y\Big). \tag{19}$$

An effective Hamiltonian defined by the exact diagonalization has the form:

$$H = \sum_{\alpha,\beta} C_{\alpha,\beta}\hat{\tau}_\alpha\hat{\sigma}_\beta \equiv C_{xo}\cdot\hat{\tau}_x\hat{\sigma}_o + C_{xy}\cdot\hat{\tau}_x\hat{\sigma}_y + C_{yx}\cdot\hat{\tau}_y\hat{\sigma}_x + H_D \ , \tag{20}$$

where H_D is diagonal in the $\hat{\tau}_z$-representation. The coefficients C_{ij} are presented in Figures 1.a, 1.b and 1.c (dashed lines) together with the corresponding analytical values from (19) (solid lines) as functions of magnetic field at $J = 40, K\perp = 20, K_\| = 1, \omega_o = 0.2, S_1 = S_2 = 10, \sigma = 1/2$. In Fig. 1.d the coefficient α extracted from C_{yx} (dashed line) is shown together with its analytical analog from (19) (solid line) as a function of sublattice spin at $J = 60, K_\perp = 20, K_\| = 1, \omega_o = 0.2, \sigma = 1/2$. This coefficient describes the interaction with the spin bath and, hence is the most interesting for the present investigation. As one can see from all the Figures, the coincidence of the analytical results with the numerical ones is quite satisfactory. Some differences between the two values for α at small S_2 are explained by the fact that semiclassical description is far from being valid in this region.

IV. CONCLUSIONS

Though the *detailed* analysis of all the factors effecting the tunneling process is beyond the scope of this paper, it is possible to make some qualitative conclusions (even without rigorous calculations with obtained Hamiltonians) on the spin dynamics of the investigated systems: (i) the quantum tunneling effect is really more strong in antiferromagnetic nanoparticles than in ferromagnetic ones that can be seen even from the fact that $A_o(0) < A_o(S_1)$; (ii) the presence of the excess spin strongly effects the quantum tunneling process due to the mere fact that it renormalizes A_o and Ω_o which both define the tunneling splitting and the bounce frequency; (iii) interaction with the environmental spins is extremely important since it can drastically change the tunneling picture down to the full suppression of it, as in the case of tunneling of half-integer excess spin or half-integer "giant spin".

ACKNOWLEDGMENTS

The author is grateful to N. Prokof'ev, P.C.E. Stamp and B. Barbara for very helpful discussions. The author also acknowledges the hospitality of the Louis Neel Laboratory (Grenoble, France), where a part of this work has been done. This work was supported by the Russian Foundation for Basic Research (grant 97-02-16548).

---

[1] J.L. Van Hammen and A. Suto, Europhys. Lett., 1, 481 (1986); E.M. Chudnovsky and L. Gunter, Phys. Rev. Lett **60**, 661 (1988); P.C.E. Stamp, E.M Chudnovsky and B. Barbara, Int. J. Mod. Phys. 6, 1355 (1992).

[2] Quantum Tunneling of Magnetization - QTM'94. edited by L. Gunter and B. Barbara, (1995).

[3] I.V. Krive and O.B. Zaslavskii, J. Phys.: Cond. Matter. 2, 9457 (1990).

[4] B. Barbara and E. M. Chudnovsky, Phys. Lett. **A**, 145, 205 (1990); E.M. Chudnovsky, p.76 in Ref.[2].

[5] E.N. Bogachek and I.V. Krive, Phys. Rev. **B46**, 14559 (1992).

[6] G. Levine and J. Howard, Phys. Rev. Lett **75**, 4142 (1995);

[7] B.A. Ivanov and A.K. Kolezhuk, cond-mat/9706292.

[8] S.E. Barnes, R. Ballou, B. Barbara and J. Strelen, Phys. Rev. Lett. **79**, 289 (1997).

[9] A. Chiolero and D. Loss, cond-mat/9708042.

[10] W. Wernsdorfer, K. Hasselbach, E. Bonet Orozco, D. Mailly, O. Kubo and B. Barbara, Phys. Rev. Lett. **79**, 4014 (1997); J. Tejada, X.X. Zhang, E. del Barco, J.M. Hernandez and E. M. Chudnovsky, Phys. Rev. Lett **79**, 1754 (1997).

[11] P.C.E. Stamp, Nature, **359**, 365 (1992): Proc. LT-20 Conf. (August, 1993; Eugene, Oregon), published in Physica **B197**,133 (1994).

[12] N.V. Prokof'ev and P.C.E. Stamp, J. Phys. CM Lett. **5**, L663-L670 (1993); N.V. Prokof'ev and P.C.E. Stamp, p.347 in Ref. [2]; N.V. Prokof'ev, P.C.E. Stamp, J. Low Temp. Phys., **104**, 143 (1996); A. Garg, Phys. Rev. Lett. **71**, 4249 (1993);

[13] D.D. Awshalom, J.F. Smyth, G. Grinstein, D.P. DiVincenzo and D. Loss, Phys. Rev. Lett. **68**, 3092 (1992).

[14] S.N. Burmistrov and L.B. Dubovskii, The applications of the quasiclassical method to the 1D problems. Preprint IAE. IAE-3881/1 (1984).

[15] I.S. Tupitsyn, N.V. Prokof'ev and P.C.E. Stamp, Int. J. Mod. Phys. **B11**, **No.24**, 2901 (1997).

[16] E. Fradkin, Field Theories of Condensed Matter Systems, (Frontiers in physics, No.82), Addison-Wesley Publishing Company, (1991).

[17] A.F. Andreev and V.I. Marchenko, Sov. Phys. Usp. **23**, 21 (1980).

[18] A. Garg, p.273 in Ref. [2].

FIGURE CAPTIONS

Figure 1 (a)-(c) The numerical coefficients C_{ij} (dashed line) in comparison with the analytical expressions from (19) (solid line) as functions of magnetic field H_y at $J = 40, K\perp = 20, K_{\|} = 1, \omega_o = 0.2, S_1 = S_2 = 10, \sigma = 1/2$. (d) The numerical (dashed line) and the analytical (19) (solid line) dependences of the coefficient α on the value of the sublattice spin at $J = 40, K\perp = 20, K_{\|} = 1, \omega_o = 0.2, \sigma = 1/2$.


Cxo
a)
2
0
-2
-4
-80
-40
0
40
80
H y
Cxy
b)
0.6
0
-0.6
-1.2
(x10-3)
-80
-40
0
40
80
H y
Cyx
c)
0.4
0
-0.4
-0.8
(x10-2)
-80
-40
0
40
80
H y
α(0)
d)
0.9
0.6
0.3
0
(x10-2)
0
4
8
12
16
S2